



The TIMI Trials

TIMI 10B

OBJECTIVE: To compare the angiographic efficacy and safety of several doses of TNK with front loaded rtPA for the treatment of ST elevation myocardial infarction.

1. Cannon CP, Gibson CM, McCabe CH, Adgey AA, Schweiger MJ, Sequeira RF, Grollier G, Giugliano RP, Frey M, Mueller HS, Steingart RM, Weaver WD, Van de Werf F, Braunwald E. TNK-tissue plasminogen activator compared with front-loaded alteplase in acute myocardial infarction: results of the TIMI 10B trial. *Circulation* 1998;98:2805-14.
2. Gibson M, McLean C, Rizzo MJ, Martin NE, Marble SJ, Dotani I, McCabe CH, Cannon CP, Braunwald E, for the TIMI 10 Investigators. Use of the TIMI frame count to compare flow at 90 min. after t-PA vs. TNK administration. *Circulation* 1996;94[Suppl. I]:I-89.
3. Gibson M, Rizzo MJ, McLean C, Sparano AM, Ryan K, Moynihan J, Ralston A, Dodge T, Cannon CP, Braunwald E. Adjunctive stenting following thrombolysis in TIMI 10A & B. *Circulation* 1997;96(Suppl. I):I-328.
4. Gibson M, Cannon CP, McCabe CH, Van de Werf FJ, Braunwald E. A randomized comparison of TPA with TNK using the TIMI Frame Count: Results of TIMI 10B. *Circulation* 1997;96(Suppl. I):I-330-I-331.
5. Goel M, Rizzo MJ, Syed J, Dodge JT, Al Mousa E, Marble SJ, Coulter SA. Relationship between proximal and distal lesion location and baseline ST-segment changes in acute myocardial infarction. *Circulation* 1998;98:I-79.
6. Gibson M, Schweiger M, Sequeira RF, Frey M, Cannon CP, Williams DO, Van de Werf F, Braunwald E, for the TIMI 10B Study Group. Outcomes of adjunctive PTCA/stenting for TIMI grade 2 flow following thrombolysis. *J Am Coll Cardiol* 1998;31(Suppl. A):231A.
7. Gibson M, Sparano A, Ryan K, Moynihan J, Kelley M, Rizzo M, Marble S, McCabe C, Dodge T, Cannon CP, TIMI 10B Study Group. Flow distal and not proximal to the lesion is a correlate of clinical outcomes after thrombolysis. *J Am Coll Cardiol* 1998;31 (Suppl A):492A.
8. Gibson M, Sparano A, Ryan K, Rizzo M, Kelley M, Marble S, Dodge T, Cannon CP, Braunwald E, for the TIMI 10B Investigators. A new angiographic method to calculate coronary artery velocity and its relationship to clinical outcomes after thrombolysis. *J Am Coll Cardiol* 1998;31 (Suppl A):12A.
9. Moynihan J, Ryan K, Sparano A, Kelly M, Rizzo M, Marble S, Cannon CP, McCabe CH, Gobson M, for the TIMI 10A & B Study Group. The range of QCA velocities for TIMI grades 2 and 3 flow. *J Am Coll Cardiol* 1998;31(Suppl. A):11A.
10. Rizzo MJ, Ryan KA, Kelley MB, Moynihan J, Cannon CP, Braunwald E. Earlier time to treatment is associated with improved coronary blood flow in acute MI. *Circulation* 1998;98 (Suppl I):I-698.
11. Gibson CM, Cannon CP, Murphy SA, Adgey J, Schweiger MJ, Sequeira RF, Grollier G, Fox NL, Weaver D, Van de Werf F, Braunwald E. Weight adjusted dosing of TNK-tissue plasminogen activator and its relationship to angiographic outcomes in the TIMI 10B Trial. *Am J Cardiol* 1999; 84:976-80.
12. Gibson CM, Murphy SA, Rizzo MJ, Ryan KA, Marble SJ, McCabe CH, Cannon CP, Van de Werf F, Braunwald E. Relationship between TIMI frame count and clinical outcomes after thrombolytic administration. Thrombolysis In Myocardial Infarction (TIMI) Study Group. *Circulation* 1999;99:1945-50.
13. Gibson CM, Ryan KA, Kelley M, Rizzo M, Mesley R, Murphy S, Swanson J, Marble SJ, Dodge JT, Cannon CP, Antman EM. Methodologic drift in the assessment of TIMI grade 3 flow and its implications with respect to reporting of angiographic trial results. *Am Heart J* 1999;137:1179-1185.
14. Tanasijevic MJ, Cannon CP, Antman EM, Wybenga DR, Fischer GA, Grudzien C, Gibson CM, Winkelman JW, Braunwald E. Myoglobin, creatine-kinase-MB and cardiac troponin-I 60-minute ratios predict infarct-related artery patency after thrombolysis for acute myocardial infarction: results from the Thrombolysis in Myocardial Infarction study (TIMI) 10B. *J Am Coll Cardiol* 1999;34:739-47.

15. Barron HV, Cannon CP, Murphy SA, Braunwald E, Gibson CM. Association between white blood cell count, epicardial blood flow, myocardial perfusion, and clinical outcomes in the setting of acute myocardial infarction: A Thrombolysis In Myocardial Infarction 10 substudy. *Circulation* 2000; 102: 2329-2334.
16. Gibson CM, Cannon CP, Murphy SA, Ryan KA, Mesley R, Marble SJ, McCabe CH, Van De Werf F, Braunwald E. Relationship of TIMI myocardial perfusion grade to mortality after administration of thrombolytic drugs. *Circulation* 2000;101:125-30.
17. Gibson CM, Anshelevich M, Murphy S, Luu L, Hynes C, Kliris J, Kermgard S, Otten MH, Antman EM, Cannon CP, Braunwald E. Impact of injections during diagnostic coronary arteriography on coronary patency in the setting of acute myocardial infarction from the TIMI trials. *Am J Cardiol.* 2000;86:1378-9.
18. Modi NB, Fox, NL, Clow FW, Tanswell P, Cannon CP, Van de Werf F, Braunwald E. Pharmacokinetics and pharmacodynamics of tenecteplase: results from a phase II study in patients with acute myocardial infarction. *J Clin Pharm* 2000; 40: 508-515.
19. Gibson CM, Marble SJ. Issues in the assessment of the safety and efficacy of tenecteplase (TNK-tPA). *Clin Cardiol* 2001;24:577-84.
20. Giugliano RP, McCabe CH, Antman EM, Cannon CP, Van de Werf F, Wilcox RG, Braunwald E. Lower-dose heparin with fibrinolysis is associated with lower rates of intracranial hemorrhage. *Am Heart J* 2001;141:742-50.
21. Schweiger MJ, Cannon CP, Murphy SA, Gibson CM, Cook JR, Antman EM, Braunwald E, for the TIMI 10B and TIMI 14 Investigators. Early coronary intervention following pharmacologic therapy for acute myocardial infarction – the combined TIMI 10B-TIMI 14 experience. *Am J. Cardiol* 2001; 88:831-6
22. Srinivas VS, Cannon CP, Gibson CM, Antman EM, Greenberg MA, Tanasijevic MJ, Murphy S, de Lemos JA, Sokol S, Braunwald E, Mueller HS. Myoglobin levels at 12 hours identify patients at low risk for 30-day mortality after thrombolysis in acute myocardial infarction: a Thrombolysis in Myocardial Infarction 10B substudy. *Am Heart J* 2001;142:29-36.
23. Gibson CM, Cannon CP, Murphy SA, Marble SJ, Barron HV, Braunwald E. Relationship of the TIMI myocardial perfusion grades, flow grades, frame count, and percutaneous coronary intervention to long-term outcomes after thrombolytic administration in acute myocardial infarction. *Circulation* 2002;105:1909-13.
24. Gibson CM, Chen M, Angeja BG, Murphy SA, Marble SJ, Barron HV, Cannon CP. Precordial ST-segment depression in inferior myocardial infarction is associated with slow flow in the non-culprit left anterior descending artery. *J Thromb Thrombolysis* 2002;13:9-12.
25. Murphy SA, Gibson CM, Van de Werf F, McCabe CH, Cannon CP. Comparison of errors in estimating weight and in dosing of single-bolus tenecteplase with tissue plasminogen activator (TIMI 10B and ASSENT I). *Am J Cardiol* 2002;90:51-4.
26. Gibson CM, Murphy SA, James D, Wong GC, Cannon CP, Giugliano RP, Antman EM, Braunwald E. Is recurrent myocardial infarction following thrombolytic administration associated with long-term mortality? *J Am Coll Cardiol* 2002;35(Suppl A):280A.